



The Heavy Metal Pollution Assessment of Topsoil in the North Bank of the Yellow River in Shandong Based on Granulation

Han Danmei^{1,2}, Zhu Lidong^{1,*}, Ding Min^{2,3,*}, Peng Shuzhen², Zhao Qiuyue², Zhang Wei²

¹College of Geography and Environment Sciences, Zhejiang Normal University, Jinhua, China

²School of Tourism, Taishan University, Tai'an, China

³Qingdao National Laboratory for Marine Science and Technology, Qingdao, China

Email address:

zhulidong@zjnu.cn (Zhu Lidong), dingmintsjy@163.com (Ding Min)

*Corresponding author

To cite this article:

Han Danmei, Zhu Lidong, Ding Min, Peng Shuzhen, Zhao Qiuyue, Zhang Wei. The Heavy Metal Pollution Assessment of Topsoil in the North Bank of the Yellow River in Shandong Based on Granulation. *International Journal of Environmental Monitoring and Analysis*. Vol. 9, No. 6, 2021, pp. 193-200. doi: 10.11648/j.ijema.20210906.15

Received: November 30, 2021; **Accepted:** December 23, 2021; **Published:** December 29, 2021

Abstract: With the rapid development of the economy, environmental pollution is becoming more and more serious. As one of the important carriers of industry and agriculture, the soil is one of the key concerns of people. The research on soil heavy metal pollution is also progressing day by day, and there are abundant research results. However, there are few studies on soil heavy metal pollution based on a granular level, most of which are whole-rock samples. Therefore, we evaluated the soil heavy metal pollution in the north bank of the Yellow River in Shandong province from the perspective of grain size. We hope to fill that gap. In this study, we systematically sampled topsoil from 15 sites on the north bank of the Yellow River in Shandong province. It was divided into two components, <20 μ m and 20-63 μ m, with a total of 32 data. The contents of heavy metals in soil were determined by ICP-MS. Based on experimental data, the single factor method and comprehensive index method were used to evaluate heavy metal pollution. The results showed that heavy metal pollution conditions were related to soil particle thickness. The contents of heavy metals in fine soil <20 μ m were higher. Among them, the heavy metal Ni content in 15 points exceeded the safety line, and 2 areas were slightly polluted. The composite index of 16 samples exceeded the safety line, and four sites had minor contamination. The heavy metal content of the soil with 20~63 μ m, thick particles was generally low, and the pollution evaluation result was clean. The results verified the geochemical characteristics of heavy metals enrichment in fine soil. In addition, the potential ecological hazards of heavy metals in soil particles < 20 μ m and 20-63 μ m were evaluated. The results showed slight ecological damage, which may be related to the biological toxicity of heavy metals.

Keywords: Shandong Province, The North of the Yellow River, Heavy Metals, Pollution Assessment, Granulation

1. Introduction

With the development of industry and agriculture, heavy metal pollution in the soil environment has become a major killer restricting the sustainable development of soil and harming the ecological environment and human health [1-3]. Soil heavy metal pollution is characterized by long residual time, strong concealment, high toxicity, small migration, etc., with strong pollution [4]. Soil heavy metal pollution and hazard assessment has always been an important research field in environmental science [5]. In the past decade, the

economy of the lower Yellow River region in Shandong province has developed rapidly, and various environmental problems, especially soil pollution, have become more and more serious. It is urgent to study soil heavy metals [6]. In addition, the accumulation, migration, and transformation mechanism of heavy metals, especially their transfer to animals and humans along the food chain, have gradually attracted public attention [2, 7].

At present, abundant achievements have been made in the

study of heavy metal content and environmental quality conditions in the soil of the Old Yellow River Channel in Shandong province, Dagu River in Qingdao, Binhai in Shandong Province, surface sediments of the Yellow River, and eastern Shandong Province [8-10]. However, there are few studies on soil heavy metal pollution assessment based on grain size in China, most of which are whole-rock samples [11]. In particular, the assessment of soil heavy metal pollution in the Shandong area of the north of the Yellow River is blank. In addition, most studies analyze and evaluate the soil heavy metal pollution in Shandong province from the micro-level, without studying the pollution situation from a specific point of view. Therefore, this study hopes to study the heavy metal content in the topsoil of the north of the Yellow River by using the method of grain grading and carrying out a quantitative quality evaluation. Here are the key parts of the job [12].

In this paper, the contents of heavy metals in the surface soil of the north bank of Yellow River in Shandong province were determined by particle size samples ($<20\mu\text{m}$ and $20\text{--}63\mu\text{m}$), and the pollution and potential ecological harm of heavy metals were evaluated. This is not only conducive to an accurate evaluation of land pollution but also conducive to research and analysis of land pollution sources. In the process of wind transport, fine components $<20\mu\text{m}$ can travel long distances in suspension up to 5KM. The crude fractions of $20\text{--}63\mu\text{m}$ were mainly proximal particles. Therefore, it is of great practical significance to conduct experimental detection and quantitative environmental quality assessment of soil heavy metals in this area [13].

2. Material and Method

2.1. Region and Sampling

The study area is located in the north of the Yellow River in central Shandong province and belongs to the North China Plain. The terrain is inclined from southwest to northeast. The climate is warm temperate semi-humid monsoon climate, with an average annual temperature of $12\text{--}15^\circ\text{C}$ and annual precipitation of $600\text{--}800\text{mm}$ [14]. The sampling points on the north bank of the Yellow River in the study area include 15 points including Donga, Chiping, Qihe, Jiyang, Huimin, Binzhou, Qingyun, Ningjin, and Dezhou (Figure 1). The sampling time was November 2015, and the sampling method was mixed sampling within 20cm from the surface. A total of 16 mixed samples were collected from each site (except Huimin Qinghe). Specific sampling sites and planting conditions are shown in Figure 1 and Table 1.




Figure 1. Illustration of sampling sites of the north of the yellow river in Shandong.

Table 1. Sampling list of north china plain and flood plain in Shandong along the Yellow River.

Sampling Site	Surface Condition	Soil Status	Sample No.
Right side of G105 of Zhaozhuang Village, Dong'e County (Yangguang Road, Jingshengzhao Village)	Wheatfield	Loess like soil	DO
Right side of G105, north of Dong'e Xiwangji Fertilizer Plant	Cotton field	Loess like soil	DOH
Next to Guanshi and 309 Qiaoxi, Chiping County	Cotton field	Brownish fragmentary, lacustrine like	CP
North of Jiaomiao, Chipping County (turning point from Chipping to Qihe County)	Open space next to the wheat field	Loess like soil	JM
Auxiliary bridge of Qihe County	Yellow Riverbank	Yellow River sand	JQHHS
South of Sangzi Town, Qihe County (Near Dingzhuang Village, the intersection of Sangxin Road and 309)	Open space	Loess like soil	SZ
Next to G220, east of Qudi Toll Station, Jiyang County	Cornfield	Loess mixed	QD
Zhaowangzhuang Village, Jiyang	Wheatfield	Loess like soil	ZWZ
Yellow River Bridge, Qikou, Qinghe Town, Huimin County	Yellow Riverbank	Low terrace, fine sand	QHHHS1
Beiyihezhuan Village, Binzhou City	Open space	High terrace, sand	QHHHS2
Right side of the highway intersection of Binzhou City	Grove	Dark soil	BZB
Cuijia Village, South External Circle, Southwest Qingyun County	Cornfield	Dark soil	BG
Next to S314, Dongrenguanzhuang Village, Ningjin County	Cornfield	Loess like soil	QYX
Next to S314, Dongweiwangzhuang Town, Dezhou City	Grove	Loess like soil	NJ
Dongnan Xiaozandian Village, Dezhou City	Wheatfield	Loess like soil	DZD
			DZN

2.2. Experiment Method

The granulation experiment is adopted with wet screening-sedimentation, and there are mainly 4 steps: (1)

take 25g naturally dried soil samples and remove the organic matter and carbonate with 30% hydrogen peroxide (H_2O_2) and 1mol/L acetic acid respectively; add 10ml 0.05mol/L (NaPO_3)₆; then put into an ultrasonic cleaner to have an

ultrasonic cleaning. (2) Remove the component of $>63\mu\text{m}$ thick particles with 230 meshes ($63\mu\text{m}$). (3) Gain the component of $<20\mu\text{m}$: the fine component screened with wet screening is transferred to the sedimentation barrel, and extract its suspension according to the Stokes law. The sedimentation time of $20\mu\text{m}$ is confirmed according to the room temperature, and it should be repeated 8-10 times till the supernatant of the upper layer is transparent. Stir quickly for 3 minutes each time and make a centrifugal concentration for the extracted suspension of $<20\mu\text{m}$ component to pour the supernatant and transfer the rest to a beaker for drying in a 40°C incubator. (4) Gain the component of $20\sim 63\mu\text{m}$: Make the centrifugal concentration for the component of $20\sim 63\mu\text{m}$ in the sedimentation barrel to pour the supernatant and transfer the rest to a beaker for drying in a 40°C incubator.

The measurement of heavy metal content: The 32 samples of $<20\mu\text{m}$ and $20\sim 63\mu\text{m}$ gained by centrifugal concentration are ground to be under 200 meshes, and take 2g respectively. The heavy metal factor, Cr, Pb, Cu, Zn, and Ni are tested in the Dyeing Control and Resource Research Laboratory of Nanjing University with an inductively coupled plasma mass spectrometer (ICP-MS).

Based on the experimental data, the study is adopted with the single factor and comprehensive pollution index method in the method of pollution index to have a pollution assessment on the heavy metal content of soil particles of $<20\mu\text{m}$ and $20\sim 63\mu\text{m}$; and it is adopted with the potential ecological hazards index method to have an evaluation on the potential ecological hazards of heavy metal.

3. The Pollution Assessment of Heavy Metal of Soil Based on Granulation

3.1. Assessment Methods and Assessment Standards

The pollution index method of single factor and comprehensive pollution index method is the common methods of the assessment of soil environment quality [14, 15]. For the pollution index method of a single factor, the pollution degree is lighter when the indexes are smaller. The assessment formula of the pollution index method of a single factor is as follow:

$$P_i = C_i / S_i$$

In the formula, P_i is the single pollution index of the pollutant I ; C_i is the measured data of pollutant I ; S_i is the assessment standard of pollutant. The soil environment is clean and safe when $P_i \leq 0.7$; and it could be called clean when $0.7 < P_i \leq 1.0$. The soil is slightly polluted when $1.0 < P_i \leq 2.0$. The soil is moderately polluted when $2.0 < P_i \leq 3.0$. The soil is seriously polluted when $P_i \geq 3.0$.

The comprehensive pollution index is adopted with N. C. Nemerow pollution index method. N. C. Nemerow reflects a comprehensive pollution degree of various pollutants on soil environmental conditions; meanwhile, it also highlights the impact of highly-concentrated pollutants on soil

environmental quality. The calculation formula is:

$$P_N = \sqrt{\frac{P_i^2 + P_i^2(\max)}{2}}$$

In the formula, P is the comprehensive pollution index of soil; P_i is the average value of all pollutants in the soil; $P_i(\max)$ is the maximal pollution index of a single pollutant in the soil. The grading standard of the method is perfectly the same as the single factor index method.

Because the soil plantation in the study area is based on the dryland in cultivated land, and fewer groves and river lands are also drylands, so the secondary standard of the National Environmental Quality Standard for Soils (announced in 1995) is selected, which means that the standard for soil environmental quality of cultivated land (dryland) is used as the standard of the heavy metal pollution assessment (Table 2) in this time [16]. The heavy metal of soil is significantly affected by the PH value. The PH measured value in the area is higher than 7.5, so the soil quality standard with the $\text{PH} > 7.5$.



3.2. The Result of Heavy Metal Content of $<20\mu\text{m}$ Fine Granular Soil and Its Pollution Assessment

The result of heavy metal content, assessment standard, and pollution assessment of $<20\mu\text{m}$ fine granular soil is shown in Table 2 and Figures 2 and 3. The heavy metal content of $<20\mu\text{m}$ fine granular soil is higher. The average content of Cr is 86.9mg/kg ; the average content of Pb is 32.3mg/kg ; the average content of Cu is 37.4mg/kg ; the average content of Zn is 111.8mg/kg and that of Ni is 51.0mg/kg . The coefficient of variation of all elements is smaller, and all of them are smaller than 0.12. The heavy metal content exceeds the background values of soil in Shandong Province. The bioconcentration factor of Ni (K_i) is 2.2, which has been 2.2 times of background values. The assessment standard of each element is based on the secondary standard of the National Environmental Quality Standard for Soils in dryland (GB15618-1995). The assessment result of the single factor pollution index shows that the heavy metal pollution degree of all places, except for South of Sangzi Town, Qihe Town (SZ), and Southwest Qingyun County (QYX) (whose pollution order is $\text{Ni} > \text{Cr} > \text{Zn} > \text{Cu} > \text{Pb}$), are $\text{Ni} > \text{Zn} > \text{Cr} > \text{Cu} > \text{Pb}$. The pollution of Ni in topsoil is the highest. The pollution index of Ni in the South of Sangzi Town, Qihe Town (SZ), and Southwest Qingyun County (QYX) are 1.08 and 1.23 respectively, which has exceeded the light pollution. While the Ni in other places is in the range of 0.7-0.98, which has exceeded the security cordon, so they should be widely focused. The comprehensive pollution index shows that the Auxiliary Bridge of Qihe County (JQHHS), South of Sangzi Town, Qihe Town (SZ), Southwest Qingyun County (QYX), and Zhaozhuang Village, Dong'e County (DO) have reached light pollution since their pollution indexes are $1.0 < P_i \leq 2.0$. Others have exceeded the security cordon to reach 0.78-0.97.

Table 2. Result of Heavy Metal Content of <20 μ m Fine Granular Soil and Pollution Assessment Result.

Location	Content (mg/Kg)					Single Factor Pollution Index (Pi)					Comprehensive Pollution Index (P _N)	Assessment Result
	Cr	Pb	Cu	Zn	Ni	Cr	Pb	Cu	Zn	Ni		
QH-HHS1	77.6	30.4	31.6	102.9	43.5	0.31	0.09	0.32	0.34	0.73	0.81	Basically clean
QH-HHS2	94.3	37.2	39.9	125.3	52.3	0.38	0.11	0.40	0.42	0.87	0.97	Basically clean
JQ-HHS	100.3	38.2	43.1	143.2	55.6	0.4	0.11	0.43	0.48	0.93	1.04	Light pollution
NJ	91.6	32.9	38.2	118.8	51.2	0.37	0.09	0.38	0.4	0.85	0.95	Basically clean
DZD	78.4	29.1	34.9	104.8	44.3	0.31	0.08	0.35	0.35	0.74	0.82	Basically clean
SZ	110.9	34.2	40.8	108.3	64.6	0.44	0.1	0.41	0.36	1.08	1.18	Light pollution
DZN	76	30.2	33.4	106.6	42	0.3	0.09	0.33	0.36	0.7	0.78	Basically clean
JM	76.9	25	28.8	100.3	43.3	0.31	0.07	0.29	0.33	0.72	0.80	Basically clean
QYX	100.1	34.3	39.4	113.2	73.7	0.4	0.1	0.39	0.38	1.23	1.33	Light pollution
ZWZ	83	26.1	32.7	101.2	43.9	0.33	0.07	0.33	0.34	0.73	0.82	Basically clean
DOH	83.9	29.5	36	105.1	48.1	0.34	0.08	0.36	0.35	0.8	0.89	Basically clean
CP	81.5	34.1	42.1	110.9	51	0.33	0.1	0.42	0.37	0.85	0.95	Basically clean
BD	75.7	31.1	38.9	105.9	48.2	0.3	0.09	0.39	0.35	0.8	0.89	Basically clean
QD	80.5	31.9	35.9	102.1	47.2	0.32	0.09	0.36	0.34	0.79	0.87	Basically clean
BZB	82.9	31.3	34.7	113.2	48.8	0.33	0.09	0.35	0.38	0.81	0.90	Basically clean
DO	96.6	41.1	48.4	126.4	58.7	0.39	0.12	0.48	0.42	0.98	1.09	Light pollution
Average Value	86.88	32.30	37.41	111.77	51.02							
Coefficient of Variation	0.10	0.10	0.10	0.08	0.12							
National Standard	250	350	100	300	60							
Background values in Shandong	65.2	24.3	21.7	61.9	23.0							
Ki	1.33	1.33	1.72	1.81	2.22							

Note: Ki is bioconcentration factors=Measured average value/background values; National standard is the soil quality standard with PH>7.5.

**Figure 2.** Single-factor pollution index situation of heavy metal of <20 μ m fine granular soil and its assessment result.**Figure 3.** Comprehensive pollution index situation of heavy metal of <20 μ m fine granular soil and its assessment result.

3.3. The Result of Heavy Metal Content of 20~63μm Fine Granular Soil and Its Pollution Assessment

The result of heavy metal content and pollution assessment of 20~63μm thick particle soil is shown in Table 3 and Figures 4 and 5. The heavy metal content of the 20~63μm is less than the fine granular soil. The average content of Cr is 53.8mg/kg; the average content of Pb is 16.1mg/kg. The average content of Cu is 13.8mg/kg; the average content of Zn is 42.8mg/kg. The average content of Ni is 21.0mg/kg. All heavy metal contents are lower than the background values of Shandong Province, and the Ki is 0.64-0.91. According to the secondary

standard of the National Environmental Quality Standard for Soils (announced in 1995), the indexes of single factor pollution and comprehensive pollution show that the heavy metal of the soil in all collection places of the area is less than 0.42, which is far away from pollution degree. Hence, it could be concluded to be in a clean status. It means that heavy metal of soil has less bioconcentration in thick particle soil. What's more, the order of heavy metal pollution of thick particle soil is different from the fine granular soil, shown to be Cr>Zn>Ni>Cu>Pb, which means that the particle size effect of each pollutant is different.

Table 3. Result of the heavy metal content of 20~63μm thick particle soil and pollution assessment result.

Location	Content (mg/Kg)					Single Factor Pollution Index (Pi)					Comprehensive Pollution Index (P _N)	Assessment Result
	Cr	Pb	Cu	Zn	Ni	Cr	Pb	Cu	Zn	Ni		
QH-HHS1	50.6	15.79	13.1	42.29	20.34	0.20	0.05	0.13	0.14	0.10	0.24	Clean
QH-HHS2	55.18	16.5	14.57	43.48	19.87	0.22	0.05	0.15	0.14	0.10	0.26	Clean
JQ-HHS	77.56	17.09	15.23	42.96	19.91	0.31	0.05	0.15	0.14	0.10	0.34	Clean
NJ	43.86	14.78	11.21	33.8	15.42	0.18	0.04	0.11	0.11	0.08	0.20	Clean
DZD	53.29	15.24	11.94	35.16	21.26	0.21	0.04	0.12	0.12	0.11	0.24	Clean
SZ	52.52	15.38	13.78	39.46	19.47	0.21	0.04	0.14	0.13	0.10	0.24	Clean
DZN	63.84	15.48	12.6	40.98	29.71	0.26	0.04	0.13	0.14	0.15	0.29	Clean
JM	41	13.04	7.32	28.02	13.86	0.16	0.04	0.07	0.09	0.07	0.19	Clean
QYX	51.62	17.43	14.65	42.71	20.36	0.21	0.05	0.15	0.14	0.10	0.24	Clean
ZWZ	59.54	14.79	12.36	40.31	20.95	0.24	0.04	0.12	0.13	0.10	0.27	Clean
DOH	45.05	13.62	10.05	33.34	15.98	0.18	0.04	0.10	0.11	0.08	0.21	Clean
CP	69.54	23.88	33.65	94.12	44.47	0.28	0.07	0.34	0.31	0.22	0.42	Clean
BD	48.14	16.33	11.7	36.6	16.48	0.19	0.05	0.12	0.12	0.08	0.22	Clean
QD	50.54	15.47	12.35	42.39	16.8	0.20	0.04	0.12	0.14	0.08	0.23	Clean
BZB	41.35	15.1	10.76	33.06	14.08	0.17	0.04	0.11	0.11	0.07	0.19	Clean
DO	57.04	17.73	16.03	56.82	26.24	0.23	0.05	0.16	0.19	0.13	0.27	Clean
Average Value	53.79	16.10	13.83	42.84	20.95							
Coefficient of Variation	0.14	0.10	0.23	0.19	0.23							
National Standard	250	350	100	300	60							
Background values in Shandong	65.2	24.3	21.7	61.9	23.0							
Ki	0.83	0.66	0.64	0.69	0.91							

Note: Ki is bioconcentration factors=Measured average value/background values; National standard is the soil quality standard with PH>7.5.




Figure 4. Single Factor Pollution Index Situation of Heavy Metal of 20~63μm Thick Particle Soil and Its Assessment Result.




Figure 5. Comprehensive Pollution Index Situation of Heavy Metal of 20–63 μ m Thick Particle Soil and Its Assessment Result.

4. Assessment of Potential Ecological Risk

4.1. Assessment Method and Assessment Standard

The paper is adopted with the potential ecological hazards index (Table 4) suggested by Sweden Hakanson [16] to have an ecological hazards assessment. This method would not only think of the ecological effect and environmental effect of heavy metal, but it also considers the environmental toxicological characteristics, which means to assess according to the toxicity levels of various heavy metals and sensitivity of organisms to heavy metal pollution. The toxicity level of

heavy metals proposed by this method is $Pb > Cu > Cr = Ni > Zn$. After the range treatment of toxicity response coefficient, it can be determined as $Pb = Cu = 5$, $Cr = Ni = 2$, $Zn = 1$. The calculation formula of the ecological hazards index is:

$$Cif = Cdi / Cir$$

$$Eir = Tir \cdot Cif$$

$$RI = \sum Eir$$

In the formula, *Cif* is a pollution index of single metal; *Cdi* is the measured value of the north bank of the Yellow River; *Cir* is the heavy metal background reference value of Shandong; *Tir* is the responding factor of biological toxicity of different metals; *Eir* is the potential ecological hazards factor of single metal; *RI* is the potential ecological hazards index of multiple metals.

To reflect the feature of the area, the heavy metal background values of Shandong are selected as a benchmark (see the following Tables 2 and 3) [17-20].

Table 4. Division of Pollution Degree of Potential Ecological Hazards Index Suggested by Hakanson.

E_i^r	Pollution Degree	RI	Hazards Level
<40	Slight	<150	Slight
40~80	Moderate	150~300	Moderate
80~160	Strong	300~600	Strong
160~320	Very strong	≥ 600	Very strong
≥ 320	Extremely strong		

Table 5. Assessment Result of Potential Ecological Hazards of Heavy Metal in Topsoil in the North Bank of the Yellow River in Shandong.

Potential Ecological Hazards Coefficient													
	<20μm						20~63μm						Assessment Result
	E ⁱ _r					RI	E ^r _r					RI	
	Cr	Pb	Cu	Zn	Ni		Cr	Pb	Cu	Zn	Ni		
QH-HHS1	2.38	6.24	7.28	1.66	3.79	21.35	1.55	3.25	3.02	0.68	1.77	10.27	Slight Ecological Hazards
QH-HHS2	2.89	7.66	9.19	2.02	4.54	26.31	1.69	3.40	3.36	0.70	1.73	10.88	
JQ-HHS	3.08	7.86	9.93	2.31	4.84	28.02	2.38	3.52	3.51	0.69	1.73	11.83	
NJ	2.81	6.77	8.79	1.92	4.45	24.75	1.35	3.04	2.58	0.55	1.34	8.86	
DZD	2.41	5.99	8.05	1.69	3.85	21.98	1.63	3.14	2.75	0.57	1.85	9.94	
SZ	3.40	7.04	9.41	1.75	5.62	27.22	1.61	3.16	3.18	0.64	1.69	10.28	
DZN	2.33	6.22	7.70	1.72	3.65	21.62	1.96	3.19	2.90	0.66	2.58	11.29	
JM	2.36	5.15	6.63	1.62	3.77	19.52	1.26	2.68	1.69	0.45	1.21	7.29	
QYX	3.07	7.05	9.07	1.83	6.40	27.42	1.58	3.59	3.38	0.69	1.77	11.01	
ZWZ	2.55	5.38	7.53	1.63	3.82	20.91	1.83	3.04	2.85	0.65	1.82	10.19	
DOH	2.57	6.07	8.29	1.70	4.18	22.82	1.38	2.80	2.32	0.54	1.39	8.43	
CP	2.50	7.02	9.69	1.79	4.44	25.43	2.13	4.91	7.75	1.52	3.87	20.19	
BD	2.32	6.40	8.95	1.71	4.19	23.57	1.48	3.36	2.70	0.59	1.43	9.56	
QD	2.47	6.57	8.26	1.65	4.10	23.05	1.55	3.18	2.85	0.68	1.46	9.72	
BZB	2.54	6.45	8.00	1.83	4.24	23.07	1.27	3.11	2.48	0.53	1.22	8.61	
DO	2.96	8.46	11.16	2.04	5.11	29.73	1.75	3.65	3.69	0.92	2.28	12.29	
Average	2.67	6.65	8.62	1.80	4.44	24.17	1.65	3.31	3.19	0.69	1.82	10.67	

4.2. Assessment Result of Potential Ecological Hazards in Soil

The assessment of potential environmental hazards in the soil of the area is shown in table 5. For the soil component of <20 μ m, the potential ecological hazards index of each heavy

metal is $Cu > Pb > Ni > Zn > Cr$; and all of them are less than 8.62. And all of them are far less than 40. The ecological hazards in the soil of 15 places are slight ecological hazards. The average value of the potential ecological hazards index of multiple metals is 24.17, which is far less than 150. It also shows that the environmental hazards in the soil of the area are slight

ecological hazards. Given the soil component of 20~63 μ m, the potential ecological hazards index of each heavy metal and the average value of potential ecological hazards index of multiple metals is smaller, which means the potential ecological hazards are extremely low. In a word, the assessment result of ecological hazards of heavy metal in the soil in the components of <20 μ m and 20~63 μ m are in slight ecological hazards, but we also still need to keep alert.

5. Conclusion

The heavy metal pollution of topsoil in the north bank of the Yellow River in Shandong is deeply affected by the thickness of soil particles. The heavy metal content of <20 μ m fine granular soil is higher, and the pollution order of the single factor is Ni>Cr>Zn>Cu>Pb. Ni has exceeded the security cordon, and even 2 places reached the light pollution. Among them, the Ni in the soil would prevent the growth and development of crops and produce some poisons to reduce the yield of crops when it is excessive; besides, it would enter animals and human bodies through the complex food chain and food web and cause harm through bioaccumulation amplification effect [21-23]. The comprehensive indexes of 15 places exceed the security cordon, and 4 of them are in light pollution, which is related to the fact that heavy metals are easily adsorbed by clay minerals and organic colloids. As for the high content of Ni and serious pollution, it should be related to the sewage discharge from surrounding industrial and mining enterprises, and attention should be paid to human activities such as fertilization and irrigation [24]. The more specific reasons would be deeply investigated and tested. The heavy metal contents of 20~63 μ m thick particle soil are lower than the fine granular soil, and the pollution assessment results are clean. Hence, it proves a geochemical characteristic that heavy metals are apt to be concentrated and accumulated in the fine granular soil, and it could provide a reference for pollution governance and ecology restoration [25]. Because of components of <20 μ m and 20~63 μ m, the assessment result of potential ecological hazards of heavy metal is the slight ecological hazards, which is consistent with the pollution assessment result of the single factor method and the comprehensive index method that some places reach the heavy metal pollution. The former mainly think about the heavy metal contents; while the latter would consider the biological toxicity besides contents [26-30].

6. Recommendations

In this study, we evaluated the heavy metal pollution in the surface soil of the north bank of the Yellow River in Shandong province. The results also indicate that the problem of heavy metals in soil in this region cannot be ignored. Therefore, we suggest that relevant departments pay attention to this problem and carry out proper treatment and recovery of heavy metal pollution. In addition, in the following study, we hope to conduct large-scale sampling and testing from a wider range,

to achieve better research results. In addition, we can also adopt more testing and evaluation methods to study soil heavy metal pollution.

References

- [1] He DM, Wang XF, Chen LJ, et al., Evaluation of Soil Heavy Metal Pollution in A Sugarcane Field in Guangxi Based on Land Accumulation Index Method and Potential Ecological Risk Index Method [J]. *Journal of Agricultural Resources and Environment*, 2014, (02): 126-131.
- [2] Liu J, Zhang M, Zhang SY, et al., Pollution Risk Evaluation of Heavy Metals in Paddy Soils in Shandong Province [J]. *Journal of Qingdao Agricultural University (National Science)*, 2019, 36 (02): 112-118+125.
- [3] Wu B, Song JM, Li XG. Environmental Geochemical Characteristics of Heavy Metals in Surface Sediments of the Yellow River Estuary [J]. *Environmental Science*, 2013, (04): 1324-1332.
- [4] Dai JR, Potential Ecological Risks of Heavy Metals in Soils of Shandong Peninsula Blue Economic Zone [J]. *Agricultural Science & Technology*, 2019, 20 (1): 42-48.
- [5] Qiu HR, Luo JZ, Zheng GH, et al., Heavy Metal Pollution Characteristics and Potential Ecological Hazards Assessment of Sediment in Southwest River Basin [J]. *Environmental Monitoring in China*, 2012, (06): 32-36.
- [6] Li HW, Shang EP, Zhang HQ, Comparative research on Spatio-temporal variability of heavy metal pollution in cultivated soils—A case study of Huang-Huai-Hai Plain and middle reaches of the Yangtze River and Jianghuai Region [J]. *China Environmental Science*, 2018, 38 (9): 3464~3473.
- [7] Cui XT, Luan WL, Niu YB, et al., Heavy Metal Pollution and Potential Ecological Hazards Assessment of urban Soil in Tangshan [J]. *Geology in China*, 2011, (05): 1379-1386.
- [8] Wu AQ. Spatial Occurrence and Ecological Risk of Heavy Metals in Coastal Soil of Shandong Province [D]. *Liaocheng University*, 2014.
- [9] Wang B. Research on Prediction of Soil Heavy Metal Pollution and Evaluation-Early Warning of Ecological Risk [D]. *Hebei Agricultural University*, 2018.
- [10] Ni ZL, Tang FB. Qu Minghua, Ding Ming, Mo Runhong. Background Values and Safety Status of Heavy Metals in Chestnut from Shandong and Hebei [J]. *Journal of Zhejiang Agricultural Sciences*, 2012, (11): 1522-1525.
- [11] Dai JR, Pang X, Yu C, et al., Study on Soil Geochemical Reference Value, Background Values and Element Bioconcentration Characteristics in Eastern Shandong Province [J]. *Geochemical*, 2011, (06): 577-587.
- [12] Ma LJ, Li XZ, Bi HB, et al., Distribution and Environmental Effects of Heavy Metals in Intertidal Sediments of Heini Bay, Shandong Province [J]. *Marine Environmental Science*, 2011, (01): 44-47.
- [13] Liu JS, Wang RQ, Dai JL, et al., Study on Soil Environmental Background Values in the Gudao District, Shandong Province of the Yellow River [J]. *Environmental Science*, 2008, (06): 1699-1704.

- [14] Liu YJ, Zhang BH, Liu ZT, et al., A Study on Distribution of Soil Heavy Metal Pollution in Weifang City Based on Geography Information System [J]. Bulletin of Soil and Water Conservation, 2016.
- [15] Xiang HM, Wan JH, Han ZZ, et al., Spatial Distribution of Heavy Metals in Soil of Dagu River Basin [J]. Hubei Agricultural Sciences, 2016, (20): 5207-5210.
- [16] GB15618-1995, Environmental Quality Standard for Soils [S].
- [17] China National Environmental Monitoring Center. Background Values of Soil Elements in China [M]. Beijing: China Environmental Science Press, 1990.330-382.
- [18] Lin YF, Lin CY, Liu XT. Improvement and case verification of potential ecological risk index assessment method for soil heavy metal contamination [J]. Soils and Crops, 2021, 10 (4): 467-473.
- [19] Zhou QX, Song YF. Restoration Principles and Methods of Polluted Soil [M]. Beijing: Science Press, 2004. 25.
- [20] Chen BQ, Sun CH. Food Pollution and Health [M]. Beijing: Chemical Industry Press, 2002. 7-9.
- [21] Wang T, Wang YP, Yang X, et al., Engineering Application Analysis of Soil Remediation Technology in Farmland Contaminated by Heavy Metals [J]. China Resources Comprehensive Utilization, 2021, 39 (11): 130-132.
- [22] Xu JH. Mathematical Methods in Modern Geography [M]. Beijing: Higher Education Press, 2002. 121-143.
- [23] Yang RM. Pollution and Monitoring of Toxic and Harmful Chemicals in the Environment [M]. Beijing: China University for Nationalities Press, 2001.65.
- [24] Feng JG, Tao X, Yu Y, et al. Study on Pollution and Pest-Free Control Technology f Apple Orchard [J]. China Fruits, 2000, (2): 9-13.
- [25] Qin PY, Wang M, Gao ZJ, et al., PANG Xu- gui. Pollution Characteristics and Ecological Risk Assessment of Soil Heavy Metals in Tengzhou [J]. Chinese Journal of Soil Science, 2018, 49 (3): 720-726.
- [26] Ma JJ, Zhang SX, Wu HP. Effects of Nickel Pollution on the Absorption and Accumulation of Mineral Nutrients in Wheat Seedlings [J]. Chinese Journal of Eco-Agriculture, 2004. 12 (3): 94-96.
- [27] Zhang YN. Prevention and Control of Heavy Metal Pollution in Agricultural Lands Research [D]. China University of Geosciences (Beijing), 2018.
- [28] Mei S, Xun W. Study on the Evaluation Index System and Standard of Arable Land Health [J]. Advances in Engineering Research, 2016. 4: 0124-0128.
- [29] Mei S, Xun W. Distribution and Risk Assessment of Soil Heavy Metals in the Main Grain Producing Area of China [J]. International Journal of Simulation Systems, Science & Technology. 2016. 9: 6.1-6.5.
- [30] Yu XJ, Cheng SC, Bai XF. Analysis on Characteristics of Various Element Concentration in Soil and Selective Cultivation under Different Pollution Sources in Some Areas in Jinan City [J]. Shandong Land and Resources, 2016, 32 (11): 49-53.